

## ΔΕΝΔΡΟΚΟΜΙΑ

### *Aleurocanthus spiniferus*

#### Ένας σημαντικός εχθρός για τα εσπεριδοειδή

##### Βιβλιογραφία

- Byrne DN & Bellows TS (1991) Whitefly biology. Annual Review of Entomology 36, 431–457. <https://doi.org/10.1146/annurev.ento.36.1.431>
- Cioffi M, Cornara D, Corrado I, Jansen MGM, Porcelli F (2013) The status of *Aleurocanthus spiniferus* from its unwanted introduction in Italy to date. Bulletin of Insectology 66 (2), 273–281.
- Clausen CP (1978) Introduced parasites and predators of arthropod pests and weeds: a world review vi, 545 pp. Agriculture Handbook No. 480. United States Department of Agriculture, Washington, USA.
- EPPO (2020) EPPO Global Database. Available at: <https://gd.eppo.int/>.
- EPPO Datasheet ALECSN, 2020.
- EPPO (2018) PM 3/85 Inspection of places of production – Vitis plants for planting. EPPO Bulletin 48, 330-349.
- Gillespie PS (2012) A review of the whitefly genus *Aleurocanthus* Quaintance & Baker (Hemiptera: Aleyrodidae) in Australia. Zootaxa 3252, 1-42.
- Gyeletshen J, Hodges A, Hodges GS (2017) Orange Spiny Whitefly, *Aleurocanthus spiniferus* (Quaintance) (Insecta: Hemiptera: Aleyrodidae). Department of Entomology and Nematology, UF/ IFAS Extension, Gainesville (USA), pp. 1-4.
- Kapantaidaki DE, Antonatos S, Kontodimas D, Milonas P, Papachristos DP (2019) Presence of the invasive whitefly *Aleurocanthus spiniferus* (Hemiptera: Aleyrodidae) in Greece. EPPO Bulletin 49(1), 127–131.
- Mifsud D, Cocquempot C, Móhlethaler R, Wilson M, Streito JC (2010) Other Hemiptera Sternorrhyncha (Aleyrodidae, Phylloxeroidea, and Psylloidea) and Hemiptera Auchenorrhyncha. Chapter 9.4. In: Roques A et al. (Eds) Alien terrestrial arthropods of Europe. BioRisk 4(1), 511-552.
- Schrader G, Camilleri M, Diakaki RMCM, Vos S (2018) Pest survey card on *Aleurocanthus spiniferus* and *Aleurocanthus woglumi*. EFSA. <https://doi.org/10.2903/sp.efsa>.

EN-1565

- USDA (1982) Pests not known to occur in the United States or of limited distribution. No. 14. Orange spiny whitefly, *Aleurocanthus spiniferus*. USDA APHIS-PPQ, Beltsville, USA.

#### Ωμύκητες του γένους

### *Phytophthora* στην Ελλάδα

#### Διάγνωση και Αντιμετώπιση

##### Βιβλιογραφία

- Drenth A. & Sendall B. (2004). Economic impact of Phytophthora diseases in southeast Asia. In: Diversity and management of Phytophthora in South East Asia. Ed. Drenth A., Guest D.I., Australian Centre for International Agricultural Research, Canberra. ACIAR Monograph 114:10-28.
- Drenth A. & Sendall B. (2001). Practical guide to detection and identification of Phytophthora. CRC for Tropical Plant Protection 1:1-41.
- Erwin D.C. & Ribeiro O.K. (1996). Phytophthora Diseases Worldwide. American Phytopathological Society, St. Paul, p.562.
- Jung T., Orlikowski L., Henricot B. et al. (2016). Widespread Phytophthora infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of Phytophthora diseases. Forest Pathology 46:134-163. DOI: 10.1111/efp.12239
- Perlerou C., Tziros G., Vetraino A. M., Diamandis S. (2010). Phytophthora cryptogea causing ink disease of *Castanea sativa* newly reported in Greece. Plant Pathology vol.59, issue 4, pages 799-799. DOI: 10.1111/j.1365-3059.2010.02268.x
- Thomidis T. (2003). The Phytophthora diseases of stone fruit trees in Greece, Archives of Phytopathology and Plant Protection 36:69-80. DOI: 10.1080/03235400310001596891
- Ελένα, K. (1999). Μύκητες του γένους Phytophthora στην Ελλάδα, είδη-ασθένειες. Τεχνικό Δελτίο αρ.13, Μπενάκειο Φυτοπαθολογικό Ινστιτούτο, σελ. 94.
- Παναγόπουλος Χ.Γ. (2007). Ασθένειες Καρποφόρων Δένδρων & Αμπέλου. Εκδόσεις Σταμούλη Α.Ε., σελ. 606.
- Παστόπουλος Σ. (2020). Φυτόφθορες ή σήψεις του λαιμού, στα οπωροφόρα. Η αναγνώριση, η πρόληψη
- και η καταπολέμηση των ωομυκήτων αυτών. [https://plantprotect.blogspot.com/2017/09/blog-post\\_9.html](https://plantprotect.blogspot.com/2017/09/blog-post_9.html)
- Τζάμος Ε.Κ. (2007). Φυτοπαθολογία, Εκδόσεις Σταμούλη Α.Ε., σελ. 557.
- Τζάμος Σ., Αντωνίου Π. & Παρασκευόπουλος Α. (2001). Η ηλιοαπολύμανση μόνη ή σε συνδυασμό με απολυμαντικά εδάφους στην αντιμετώπιση εδαφογενών παθογόνων, μείωση της διάρκειας εφαρμογής της με αδιαπέρατα πλαστικά φύλλα. Περιοδικό Γεωργία - Κτηνοτροφία, τεύχος 4, σελ. 31-35.

και η καταπολέμηση των ωομυκήτων αυτών. [https://plantprotect.blogspot.com/2017/09/blog-post\\_9.html](https://plantprotect.blogspot.com/2017/09/blog-post_9.html)

– Τζάμος Ε.Κ. (2007). Φυτοπαθολογία, Εκδόσεις Σταμούλη Α.Ε., σελ. 557.

– Τζάμος Σ., Αντωνίου Π. & Παρασκευόπουλος Α. (2001). Η ηλιοαπολύμανση μόνη ή σε συνδυασμό με απολυμαντικά εδάφους στην αντιμετώπιση εδαφογενών παθογόνων, μείωση της διάρκειας εφαρμογής της με αδιαπέρατα πλαστικά φύλλα. Περιοδικό Γεωργία - Κτηνοτροφία, τεύχος 4, σελ. 31-35.

#### Επίδραση άρδευσης στα ποιοτικά χαρακτηριστικά ελαιόλαδου και ελαιόκαρπου

#### Θα πρέπει να είναι στοχευμένη και ορθολογική

##### Βιβλιογραφία

- Stefanoudaki E, Williams M, Chartzoulakis K, Harwood J. Olive oil qualitative parameters after orchard irrigation with saline water. J Agric Food Chem. 2009a Feb 25;57(4):1421-5. doi: 10.1021/jf8030327. PMID: 19173601.
- Stefanoudaki E, Williams M, Chartzoulakis K, Harwood J. Effect of irrigation on quality attributes of olive oil. J Agric Food Chem. 2009b Aug 12;57(15):7048-55. doi: 10.1021/jf900862w. PMID: 19722583.
- Koseoglu O, Kaya U, Nergiz C, Camoglu G, Akkuzu E, Asik S, Avci M, (2011). THE EFFECT OF DIFFERENT IRRIGATION LEVELS ON THE QUALITY OF OLIVE OIL IN THE WESTERN COASTAL REGION IN TURKEY. Acta Hortic. 2011. 888, 73-80. doi.org/10.17660/ActaHortic.2011.888.7
- Pliakoni, E. D., & Nanos, G. D. (2011). Influence of deficit irrigation and reflective mulch on “Konservolea” olive leaf physiology during the growing period. Acta Horticulturae, 888, 199– 204. <https://doi.org/10.17660/ActaHortic.2011.888.22>
- IOC (2022). World Olive Oil and Table Olive Figures. Retrieved January 31, 2022, from International Olive Council website: <https://www.internationaloliveoil.org/what-wedo/economic-affairs-promotion-unit/#figures>
- Girón, I.F.; Corell, M.; Galindo, A.; Torrecillas, E.; Morales, D.; Dell'Amico, J.; Torrecillas, A.; Moreno, F.; Moriana, A. Changes in



the physiological response between leaves and fruits during a moderate water stress in table olive trees. *Agric. Water Manag.* 2015, 148, 280–286

- Molina-Moral, J.C.; Moriana-Elvira, A.; Pirez-Latorre, F.J. The Sustainability of Irrigation Strategies in Traditional Olive Orchards. *Agronomy* 2022, 12, 64. <https://doi.org/10.3390/agronomy12010064>

## ΦΥΤΑ ΜΕΓΑΛΗΣ ΚΑΛΛΙΕΡΓΕΙΑΣ

### Πράσινο και ρόδινο σκουλήνκι στο βαμβάκι

#### Δύο εχθροί με οικονομική σημασία

##### Βιβλιογραφία

- Aheer GM, Ali A, Akram M (2009) Effect of weather factors on populations of *Helicoverpa armigera* moths at cotton-based agro-ecological sites. *Entomological Research* 29, 36–42.
- CABI Compendium, 2021. *Helicoverpa armigera* (cotton bollworm). DOI: <https://doi.org/10.1079/cabicompendium.26757>.
- Daly, 1988. Insecticide resistance in *Heliothis armigera* in Australia. *Pesticide Science* 23, Issue 2, 165–176.
- Gina E. Dillon and Gary P. Fitt, 1995. Reassessment of sampling relationships for *Helicoverpa* spp. (Lepidoptera: Noctuidae) in Australian cotton. *Bulletin of Entomological Research* 85, 321–329.
- EPPO, 1981. Data sheets on quarantine organisms No. 110, *Helicoverpa armigera*. *Bulletin OEPP/EPPO Bulletin* 11(1), 1–6.
- Gouge DH, Lee LL, Henneberry TJ, 1999. Parasitism of diapausing pink bollworm *Pectinophora gossypiella* (Lepidoptera: Gelechiidae) larvae by entomopathogenic nematodes (Nematoda: Steinernematidae, Heterorhabditidae). *Crop Protection*, 18(8):531–537; 44 ref.
- Greathead DJ, 1989. The potential of biological control in integrated pest management of insect pests in cotton. In: Green MB, Lyon DJ de B. eds. *Pest Management in Cotton*. Chichester, UK: Ellis Harwood.
- Green MB, Lyon DJ de B(Editors), 1989. Pest management in cotton. Chichester, West Sussex, UK; Ellis Horwood Limited, 259 pp.
- Hardwick DF (1965) The corn

earworm complex. *Memoirs of the Entomological Society of Canada* 40, 1–247.

- Henneberry TJ, Forlow Jech L, Burke RA, 1996. Pink bollworm adult and larval susceptibility to steinernematid nematodes and nematode persistence in the soil laboratory and field tests in Arizona. *Southwestern Entomologist*, 21(4):357–368; 20 ref.
- S. V. Leontopoulos, I. K. Vagelas and F. T. Gravanis, 2011. Estimate the Emergence of *Pectinophora gossypiella* Saunders. (Lepidoptera: Gelechiidae) with Degree Days in the Region of Thessaly, Greece. *Journal of Agricultural Science and Technology A* 1 (20 II) 182–186.
- Mabbett T, 1991. Pink bollworm management with pheromones. *International Pest Control*, 33(3):59.
- National Cotton Council of America. The story of cotton. <https://www.cotton.org/pubs/cottoncounts/story/>
- Oerke EC, Dehne HW, Schöpbeck F, Weber A, 1994. Crop production and crop protection: estimated losses in major food and cash crops. *Crop production and crop protection: estimated losses in major food and cash crops.*, xxii + 808 pp.; [ref. at ends of chapters, available from publishers at PO Box 1991, Amsterdam, Netherlands or PO Box 945, Madison Square Station, NY 10160-0757, USA].
- Pirez-Guerrero, S., Gelan-Begna, A., & Vargas-Osuna, E. (2013). Impact of *Cheiracanthium pelasgicum* (Araneae: Miturgidae) and *Chrysoperla carnea* (Neuroptera: Chrysopidae) intraguild predation on the potential control of cotton pest *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Biocontrol Science and Technology*, 24(2), 216–228. <https://doi.org/10.1080/09583157.2013.855167>
- Pogue MG (2004) A new synonym of *Helicoverpa zea* (Boddie) and differentiation of adult males of *H. zea* and *H. armigera* (Hóbner) (Lepidoptera: Noctuidae: Heliothinae). *Annals of the Entomological Society of America* 97(6), 1222–1226.
- Rajapakse CNK, Walter GH (2007) Polyphagy and primary host plants: oviposition preference versus larval performance in the lepidopteran pest *Helicoverpa armigera*. *Arthropod-Plant Interactions* 1, 17–26. Behere et al., 2013.
- Reddy GVP, Manjunatha M (2000) Laboratory and field studies on the
- integrated pest management of *Helicoverpa armigera* (Hóbner) in cotton, based on pheromone trap catch threshold level. *Journal of Applied Entomology* 124, 213–221.
- Shah KD, Jhala RC, Dhande SR (2017) Standardization of pheromone traps for the mass trapping of *Helicoverpa armigera* (Hóbner) Hardwick in tomato. *Current Agriculture Research Journal* 5, 45–49.
- MP Zalucki, G Daglish, S Firempong and P Twine, 1986. The Biology and Ecology of *Heliothis-Armigera* (Hubner) and *Heliothis-Punctigera* Wallengren (Lepidoptera, Noctuidae) in Australia - What Do We Know. *Australian Journal of Zoology* 34(6) 779–814.

## Ευρωπαϊκό έργο GOOD-Agroecology for Weeds

### Αξιολόγηση αγροοικολογικών προσεγγίσεων και στρατηγικών στο πλαίσιο της ολοκληρωμένης διαχείρισης των ζιζανίων

##### Βιβλιογραφία

- Bamal, D., Duhan, A., Pal, A., Beniwal, R. K., Kumawat, P., Dhande, S., ... & Yadav, R. (2024). Herbicide risks to non-target species and the environment: A review. *Environmental Chemistry Letters*, 1–56.
- Chauhan, B. S. (2020). Grand challenges in weed management. *Frontiers in Agronomy*, 1, 3.
- Heap, I. The International Herbicide-Resistant Weed Database. Online. Tuesday, September 10, 2024. Available [www.weedscience.org](http://www.weedscience.org)
- Mesnage, R., Szikacs, A., & Zaller, J. G. (2021). Herbicides: Brief history, agricultural use, and potential alternatives for weed control. In *Herbicides* (pp. 1–20). Elsevier.
- Ranaldo, M., Carlesi, S., Costanzo, A., & Bürberi, P. (2020). Functional diversity of cover crop mixtures enhances biomass yield and weed suppression in a Mediterranean agroecosystem. *Weed Research*, 60(1), 96–108.
- Smith, R. G., Warren, N. D., & Cordeau, S. (2020). Are cover crop mixtures better at suppressing weeds than cover crop monocultures? *Weed Science*, 68(2), 186–194.
- Travlos, I., Gazoulis, I., Simić,

M., & Kanatas, P. (2023). The Underestimated Role of Cultural Practices in Ecologically Based Weed Management Approaches. *Ecologically-Based Weed Management: Concepts, Challenges, and Limitations*, 75-92.

## ΚΤΗΝΟΤΡΟΦΙΑ

### Η σημασία της φυτοποικιλότητας

#### Στην επιλογή δίαιτας των βοσκόντων ζώων και στο περιβάλλον

##### Βιβλιογραφία

- Provenza F.D. and T.G. Papachristou, 2009. Behavior-based management of ecosystems. In: T.G. Papachristou, Z.M. Parissi (Editors), Nutritional and Foraging Ecology of Sheep and Goats. 12th Seminar of the FAO-CIHEAM Sub-Network on Sheep and Goat Nutrition, Thessaloniki, Greece, October 11-13, 2007. Options Méditerranéennes (Series A: Mediterranean seminars), 85: 13-28.
- Δημοσιεύσεις του προγράμματος BEHAVE (Behavioral Education for Human, Animal, Vegetation and Ecosystem management) που συντονίζονταν από τον Καθηγητή Frederick Provenza (Utah State University) και ο συγγραφέας ήταν μέλος της ερευνητικής ομάδας (υπότροφος του Ιδρύματος Fulbright).
- Παπαχρήστου, Θ.Γ., 2019. Τοξικά φυτά για τα ζώα: Άλλοι επίδραση επιβλαβών φυτών λιβαδιών και ζώων κατά τη βόσκηση. Γεωργία & Κτηνοτροφία, 1: 75-80.
- Παπαχρήστου, Θ.Γ., 2019. Διασύνδεση βιοχημικών συστατικών φυτών και συμπεριφοράς βόσκουσης: Συνέπειες για τα λιβάδια και τα ζώα. Πρακτικά 7ου Πανελλήνιου Συνεδρίου Τεχνολογίας Ζωικής Παραγωγής, Θεσσαλονίκη 1 Φεβρουαρίου 2019. <http://www.ap.teithe.gr/>, σελ., 47-55. ■